

hep-th/9204026

IASSNS-HEP-92/22

March 1992

Dilaton Gravity and No-Hair Theorem in Two Dimensions[★]

OLAF LECHTENFELD[†]

and

CHIARA NAPPI[‡]

*School of Natural Sciences
Institute for Advanced Study
Princeton, NJ 08540*

ABSTRACT

We study a general class of two-dimensional theories of the dilaton-gravity type inspired by string theory and show that they admit charged multiple-horizon black holes. These solutions are proved to satisfy scalar no-hair theorems.

★ Research supported in part by the Department of Energy, contract DE-FG02-90ER40542, and by the Ambrose Monell Foundation

† e-mail: OLAF@IASSNS.BITNET or LECHTENF@GUINNESS.IAS.EDU

‡ e-mail: NAPPI@IASSNS.BITNET

Over the years many attempts have been made to formulate a non-trivial theory of two-dimensional gravity (see, for example, ref. [1]). More recently, attention has been drawn to two-dimensional string-inspired models of gravity, which have been shown to have black hole solutions [2]. The non-triviality of these models arises from the non-minimal coupling of a scalar field, the “dilaton”, to the scalar curvature. In the presence of a dilaton potential of the type produced by string loop corrections, the black hole solutions may exhibit multiple horizons, as discussed in [3], where charged black holes in “heterotic” two-dimensional string theories were also studied. In this letter we put together all these features and examine a general class of two-dimensional string-inspired theories. At the classical level,[§] they all seem to provide a consistent definition of two-dimensional gravity that has many analogies with four-dimensional gravity. We exploit these analogies to prove that generalized Reissner-Nordström black-holes satisfy no-hair theorems. This puts constraints on additional scalar fields if they are to enter non-trivially in such solutions.

In the following we consider two-dimensional “dilaton gravity” conformally coupled to Maxwell and scalar fields,

$$S_2 = \int d^2x \sqrt{G} e^{-2\phi} \left[R + \gamma(\nabla\phi)^2 - \frac{1}{4}e^{\epsilon\phi}F^2 + \delta(\nabla\psi)^2 + V(\phi, \psi) \right] \quad (1)$$

where $\sqrt{G} \equiv \sqrt{-\det g_{\mu\nu}}$, ϕ denotes the dilaton, and $F_{\mu\nu} = \partial_{[\mu}A_{\nu]}$ is the Maxwell field. The scalar ψ is a spectator field, and the restriction to a single scalar is merely for simplicity. S_2 covers a large family of non-trivial actions for two-dimensional gravity. For the special choice of $\gamma = 4$, $F = \psi = 0$, and $V(\phi, 0) = -\Lambda$, this is the usual bosonic string spacetime effective action [5], discussed in [2,3]. In addition, the Maxwell term (for $\epsilon = 0$) appears for the heterotic string. The cosmological constant Λ originates from the conformal anomaly.

[§] For recent work on the quantum level, see [4].

In another specific case, $\gamma=2$, $\epsilon=0$, and $V(\phi, \psi)=2e^{2\phi}+U(\psi)$, our action is identical to the *four*-dimensional

$$S_4 = \int d^4x \sqrt{{}^{(4)}G} \left[{}^{(4)}R - \frac{1}{4}F^2 + \delta(\nabla\psi)^2 + U(\psi) \right] \quad (2)$$

for spherically symmetric and purely electrically charged field configurations, as is obvious in the gauge

$${}^{(4)}ds^2 = {}^{(2)}ds^2 + e^{-2\phi(r)}d\Omega \quad , \quad F = f(r) dr \wedge dt \quad . \quad (3)$$

Here the “dilaton” ϕ has become part of the four-dimensional metric. Four-dimensional purely *magnetic* Maxwell fields, $F = q \sin\theta d\theta \wedge d\phi$, simply reduce to a modification $V \rightarrow V - 2q^2 e^{4\phi}$ of the potential $V(\phi, \psi)$ in two dimensions. For example, all solutions of [6] can be obtained from our action (1) with $\gamma=2$, $\delta=-2$, $F=0$ and $V(\phi, \psi)=2e^{2\phi}-2q^2e^{4\phi-2a\psi}$.

Certain choices of the coefficients γ, δ, ϵ and the potential V in S_2 are related by Weyl rescaling. Of course, conformal rescaling cannot remove the direct gravity-dilaton coupling in S_2 , a unique feature of two-dimensional gravity. However, $ds^2 \rightarrow e^{\beta\phi}ds^2$ does change the coefficients in S_2 ,

$$\gamma \rightarrow \gamma - \beta^2 \quad , \quad \delta \rightarrow \delta \quad , \quad \epsilon \rightarrow \epsilon - \beta \quad , \quad V \rightarrow e^{\beta\phi}V \quad , \quad (4)$$

and can eliminate the dilaton kinetic term for $\beta = \pm\sqrt{\gamma}$. In particular, for the traditional case of $\gamma=4$ and $V = -\Lambda + U$, $\beta=2$ yields

$$S_2^{\text{trad}} \rightarrow \int d^2x \sqrt{G} \left[e^{-2\phi}R - \Lambda - \frac{1}{4}e^{(\epsilon-4)\phi}F^2 + \delta e^{-2\phi}(\nabla\psi)^2 + U(\psi) \right] \quad . \quad (5)$$

The classical spacetime generated by this action ($\gamma=0$) in the absence of Maxwell and scalar fields is flat, but the dilaton solution still carries information about the black hole singularity. Had we chosen $V = -\Lambda e^{-2\phi} + U(\psi)$ instead, another well-known candidate action for two-dimensional gravity, $\tilde{S}_2 = \int d^2x \sqrt{G} e^{-2\phi} [R - \Lambda + \dots]$,

would have emerged [1]. The transformation (4) implies that the distinction between metric and dilaton is largely an arbitrary one, since rescaling shifts structure from one to the other. Despite the simplicity of choosing $\gamma = 0$, the analogy between two-dimensional “dilaton gravity” and four-dimensional pure gravity is best pursued for nonzero γ . The similarity between the two theories will become even more evident in the gauge we will soon adopt.

The equations of motion for the general action S_2 are

$$\begin{aligned}
0 &= -2\nabla^2\phi + 4(\nabla\phi)^2 - V(\phi, \psi) - \frac{1}{4}e^{\epsilon\phi}F^2 \\
0 &= R_{\mu\nu} + 2\nabla_\mu\nabla_\nu\phi + (\gamma-4)(\nabla_\mu\phi\nabla_\nu\phi - g_{\mu\nu}(\nabla\phi)^2 + \frac{1}{2}g_{\mu\nu}\nabla^2\phi) \\
&\quad - \frac{1}{2}e^{\epsilon\phi}F_\mu^\rho F_{\nu\rho} + \frac{\epsilon}{16}g_{\mu\nu}e^{\epsilon\phi}F^2 + \delta\nabla_\mu\psi\nabla_\nu\psi - \frac{1}{4}g_{\mu\nu}\frac{\partial V}{\partial\phi} \\
0 &= \nabla^\mu(F_{\mu\nu}e^{(\epsilon-2)\phi}) \\
0 &= -2\delta\nabla^2\psi + 4\delta\nabla\phi\cdot\nabla\psi + \frac{\partial V}{\partial\psi}
\end{aligned} \tag{6}$$

where appropriate linear combinations have been formed. Our interest focuses on static classical configurations $\{F = f(r)dr \wedge dt, \phi(r), \psi(r)\}$ with asymptotically flat metric,

$$\begin{aligned}
ds^2 &= -g(r)dt^2 + g(r)^{-1}dr^2 \\
&\rightarrow -dt^2 + dr^2 \quad \text{as } r \rightarrow \infty
\end{aligned} \tag{7}$$

in the half-space $\mathbf{R}\times\mathbf{R}_+ \ni \{t, r\}$.[★] Horizons $r=h_i$ are defined as zeros of $g(r)$. Note that our choice of gauge is not the conformal one but sets $\sqrt{G} = 1$ everywhere.[†]

★ Actually, this asymptotic behavior is too restrictive: In two dimensions, a *linear* function $g(r)$ can always be gauged to $g = 1$ by a suitable coordinate transformation since $R = -g''$. Obviously, we should admit $g(r) \rightarrow A + Br$ as $r \rightarrow \infty$. In contrast, a quadratic limit, $g \rightarrow A + Br - \Lambda r^2$, signifies a nonzero cosmological constant (when $\gamma \neq 4$) and for $\Lambda > 0$ leads to a cosmological event horizon. We do not consider such theories here [7].

† For a treatment of S_2 (without the Maxwell field) in conformal gauge see [8].

The field variations (6) in gauge (7) become (primes mean $\frac{d}{dr}$)

$$\begin{aligned}
(g\phi')' - 2g\phi'^2 - \frac{1}{4}f^2e^{\epsilon\phi} + \frac{1}{2}V(\phi, \psi) &= 0 \\
2\phi'' + (\gamma-4)\phi'^2 + \delta\psi'^2 &= 0 \\
(f e^{(\epsilon-2)\phi})' &= 0 \\
(g\psi')' - 2g\phi'\psi' - \frac{1}{2\delta}\frac{\partial V}{\partial\psi}(\phi, \psi) &= 0
\end{aligned} \tag{8}$$

which already solves the Maxwell field, $f(r) = f_0 e^{(2-\epsilon)\phi(r)}$.

An analytic solution of (8) is possible for a constant spectator field, $\psi(r) = \psi_0$, sitting at a potential minimum, $\frac{\partial V}{\partial\psi}(\phi, \psi_0) = 0$. We abbreviate $V(\phi, \psi_0) =: V_0(\phi)$. With this simplification we explicitly find

$$\phi(r) = \begin{cases} \phi_0 - \frac{Q}{2}r & \text{for } \gamma = 4, \\ \phi_0 + \frac{2}{\gamma-4}\ln(r-r_0) & \text{for } \gamma \neq 4. \end{cases} \tag{9}$$

Q , r_0 and ϕ_0 are integration constants; we shall set $r_0 = 0$ without loss of generality. The remaining equation results in

$$g(r) = \begin{cases} e^{-Qr} \left[-2m + \frac{1}{Q} \int^r dr' e^{Qr'} W(\phi(r')) \right] & \text{for } \gamma = 4, \\ r^{\alpha+1} \left[-2m - \frac{1}{\alpha} \int^r dr' r'^{-\alpha} W(\phi(r')) \right] & \text{for } \gamma \neq 4, \end{cases} \tag{10}$$

with $\alpha \equiv \frac{4}{\gamma-4}$ and $W(\phi) \equiv V_0(\phi) - \frac{1}{2}f_0^2 e^{(4-2\epsilon)\phi}$. The integration constant m turns out to be the mass of the black hole [2,3]. Demanding $g(r \rightarrow \infty) \rightarrow 1$ restricts $\gamma \leq 4$ ($\alpha \leq 0$), $\epsilon \leq \gamma/2$, and forces the dilaton potential to decay (in r) as $W(\phi(r \rightarrow \infty)) \rightarrow \alpha(\alpha+1)e^{(4-\gamma)(\phi-\phi_0)}$. Obviously, a cosmological constant is allowed only for $\gamma=4$ where $W(\phi(r \rightarrow \infty)) \rightarrow Q^2 = -\Lambda$ is mandatory.

The type of dilaton potential to be expected from closed string loop corrections is of the form

$$V_{0,\text{string}}(\phi) = \sum_{n \geq 0} a_n e^{2n\phi} \quad . \tag{11}$$

Plugging this into eq. (10), one gets (setting $\epsilon=0$ for simplicity)

$$g(r) = \begin{cases} \frac{b_0}{Q^2} - 2me^{-Qr} + \frac{b_1}{Q}re^{-Qr} + \sum_{n \geq 2} \frac{b_n}{(1-n)Q^2} e^{-nQr} & \text{for } \gamma = 4, \\ -2mr^{1+\alpha} - \sum_{n \geq 0} \frac{b_n}{n\alpha^2 + \alpha(1-\alpha)} r^{2+n\alpha} & \text{for } \gamma \neq 4, \end{cases} \quad (12)$$

with $b_n = e^{2n\phi_0}(a_n - \frac{1}{2}\delta_{n,2}f_0^2)$. In case the above denominator vanishes, $n\alpha - \alpha + 1 = 0$, the contribution to $g(r)$ is $-\frac{b_n}{\alpha}r^{1+\alpha} \ln r$ instead of being singular. Asymptotic flatness requires that $\gamma = 4 - 2k$ ($\alpha = -\frac{2}{k}$) with $k = 0, 1, 2, \dots$, and $b_0, b_1, \dots, b_{k-1} = 0$ with the possible exception of $b_{k/2}$, but b_k must not vanish except for $\gamma = 0$. Furthermore, we need $m = 0$ unless $\gamma = 4, 2, 0$. These metrics generalize the well-known Reissner-Nordström black hole, as is most evident for $\gamma = 2$ ($b_0 = 0, b_1 = 2$),

$$g(r) = 1 - 2mr^{-1} - \frac{b_2}{2}r^{-2} - \frac{b_3}{6}r^{-4} - \dots \quad (13)$$

The emergence of multiple-horizon spacetimes only depends on the values of the coefficients a_n , which should be determined from string theory. If we admit a non-zero cosmological constant, $\Lambda = -a_0 = -b_0 \neq 0$, the metric function $g(r)$ for $\gamma \neq 4$ will pick up an additional term of $-\Lambda r^2$ which leads to a cosmological event horizon for $\Lambda > 0$. This is also true for the four-dimensional de Sitter gravity (2) which admits up to four horizons [7] even though 4-d general covariance restricts $V_0 = -\Lambda + 2e^{2\phi}$ in the equivalent two-dimensional action. For the two-dimensional case, examples of interesting spacetime geometries associated with multiple horizons have been discussed in [3]. Here we have generalized such solutions to the whole class (1) of dilaton-gravity-Maxwell theories.

The second issue addressed in this note is the question of *no-hair theorems*. In our point of view, the dilaton is integral part of two-dimensional gravity since it is necessary for the existence of black holes in the first place. Therefore such theorems should not apply to the dilaton [9]. The question is rather whether our solutions to the dilaton-gravity-Maxwell system can be perturbed by a non-trivial static spectator field, $\psi' \neq 0$. We know that the *four*-dimensional Einstein-Maxwell

theory obeys no-hair theorems. Given the analogy between 2-d dilaton gravity and 4-d pure gravity, it is natural to ask whether our solutions (12) are also stabilized by such theorems. If the direct spectator-dilaton coupling in (1) was absent this question would obviously be answered in the affirmative. Due to the exponential coupling, however, the matter is less than trivial. The main scope of this letter is to establish conditions for the uniqueness of the above solutions to two-dimensional “dilaton gravity” in the presence of Maxwell and scalar fields.

The key observation is that the equation of motion for ψ is the same as in four dimensions, since ψ couples in S_2 in exactly the same way as in S_4 for a spherically symmetric setting. This will allow us to take over Bekenstein’s proof of the scalar no-hair theorem [10]: The equation of motion for ψ , last of eqs. (8), can be rewritten as

$$(g e^{-2\phi} \psi')' = \frac{1}{2\delta} e^{-2\phi} \frac{\partial V}{\partial \psi}(\phi, \psi) \quad . \quad (14)$$

We multiply (14) by $(\psi - \psi_0)$ and integrate between the last horizon h , where $g(h)=0$, and spatial infinity, where $g \rightarrow 1$. Integration by parts then leads to

$$\int_h^\infty [(\psi - \psi_0) g e^{-2\phi} \psi']' = \int_h^\infty g e^{-2\phi} \psi'^2 + \frac{1}{2\delta} \int_h^\infty (\psi - \psi_0) e^{-2\phi} \frac{\partial V}{\partial \psi}(\phi, \psi) \quad . \quad (15)$$

If the boundary term vanishes and both integrands on the right hand side are non-negative, we may conclude that $\psi' \equiv 0$ and thus $\psi \equiv \psi_0$. For this to be true, however, four conditions have to be met:

- a) $\delta > 0$,[★]
- b) regularity of ψ and ϕ at $r=h$,[†]
- c) $\frac{\partial^2 V}{\partial \psi^2} \geq 0$ in $[h, \infty]$ for any ϕ ,
- d) $\psi \rightarrow \psi_0$ “fast enough” as $r \rightarrow \infty$.

★ The solutions of [6] violate this condition.

† For a violation of this condition see [11].

The conditions above are the same as in four-dimensional gravity. “Morally” speaking, the charged and multiple-horizon solutions to the general class of theories in (1) satisfy the classical no-hair theorem. In practice, however, the condition d) above is more or less restrictive depending on the asymptotic behaviour of the dilaton, *i.e.* on the value of the coefficient γ . It ensures the vanishing of the boundary term and means that $e^{-2\phi}\psi'^2$ must decay faster than $1/r$. Apparently, the fall-off assumption d) is very restrictive for the traditional dilaton case ($\gamma=4$). To get an idea, one might try to neglect the back reaction of ψ on our above solutions for ϕ and estimate that ψ should approach ψ_0 faster than $e^{-\frac{1}{2}Qr}/\sqrt{r}$ or $1/r^{1-\alpha/2}$ for $\gamma=4$ or $\gamma\neq 4$, respectively.

Fortunately, we can employ a different strategy to substantially strengthen the no-hair theorem for the *traditional* dilaton-gravity-Maxwell system. When $\gamma=4$ and $\epsilon=0$ the gravitational and dilaton equations of motion imply[‡]

$$\begin{aligned}(g\phi')' - 2g\phi'^2 &= \frac{1}{4}f^2 - \frac{1}{2}V(\phi, \psi) \\ g'' - 2\phi'g' &= f^2 - \frac{1}{2}\frac{\partial V}{\partial \phi}(\phi, \psi) \\ \phi'' &= -\frac{1}{2}\delta\psi'^2 \leq 0\end{aligned}\tag{16}$$

while the Maxwell solution is unchanged, $f = f_0 e^{2\phi}$. Since the dilaton exponential in (15) is responsible for the weakness of the standard argument, we trivially rewrite eq. (15) as

$$\int_h^\infty [(\psi-\psi_0)g\psi']' = \int_h^\infty g\psi'^2 + \frac{1}{2\delta}\int_h^\infty (\psi-\psi_0)\frac{\partial V}{\partial \psi}(\phi, \psi) + 2\int_h^\infty (\psi-\psi_0)g\phi'\psi' \quad . \tag{17}$$

The no-hair theorem may now be valid under the same conditions as above, except for a much weaker fall-off constraint, namely d') $\psi \rightarrow \psi_0$ in an arbitrary fashion at large r . However, success hinges on the positivity of the last term in (17). A

[‡] The second of these equations is a linear combination of eqs. (8) and their derivatives.

second partial integration produces

$$2 \int_h^\infty (\psi - \psi_0) g \phi' \psi' = \int_h^\infty [(\psi - \psi_0)^2 g \phi']' - \int_h^\infty (\psi - \psi_0)^2 (g \phi')' \quad (18)$$

which invites the application of the other classical equations, (16),

$$-(g \phi')' = \frac{1}{2} \delta g \psi'^2 - g' \phi' = \frac{1}{2} \delta g \psi'^2 - \frac{1}{2} g'' + \frac{1}{2} f_0^2 e^{4\phi} - \frac{1}{4} \frac{\partial V}{\partial \phi} \quad . \quad (19)$$

To proceed, obviously some information about sign and magnitude of g' and ϕ' in $[h, \infty]$ is needed. To this end we restrict ourselves again to $V(\phi, \psi) = Q^2 + U(\psi)$, *i.e.* string tree level dilaton potential. The first two of eqs. (16) can be turned into integral representations for ϕ' and g' ,

$$\begin{aligned} \phi'(r) &= \frac{e^{2\phi(r)}}{2g(r)} \int_h^r dr' \left[\frac{1}{2} f_0^2 e^{2\phi(r')} - e^{-2\phi(r')} (U(\psi(r')) + Q^2) \right] \\ g'(r) &= e^{2\phi(r)} \left[c - f_0^2 \int_r^\infty dr' e^{2\phi(r')} \right] \quad , \end{aligned} \quad (20)$$

with $c = 2mQ > 0$ from the large- r limit. The electrical charge q of the black hole hides in $\frac{1}{2} f_0^2 = q^2 Q^2$. Our asymptotic condition on g then translates to

$$1 = \int_h^\infty dr g'(r) = 2mQ \int_h^\infty dr e^{2\phi(r)} - 2q^2 Q^2 \int_h^\infty dr e^{2\phi(r)} \int_r^\infty dr' e^{2\phi(r')} \quad (21)$$

which is a quadratic equation for $\int_h^\infty e^{2\phi}$ and solved by

$$\int_h^\infty dr e^{2\phi(r)} = \frac{1}{q^2 Q} [m - \sqrt{m^2 - q^2}] \quad . \quad (22)$$

Plugging this result back into eq. (20) we can bound g' by

$$g'(r) \geq e^{2\phi(r)} 2Q \sqrt{m^2 - q^2} \geq 0 \quad (23)$$

outside the horizon! We see that g' may just vanish at the horizon in the extremal

case, $|q|=m$. The sign of ϕ' , on the other hand, need not always be negative. From its large- r behavior and the fact that $\phi'' \leq 0$ we conclude that ϕ' is maximal for minimal r , which on $[h, \infty]$ means $r=h$. Applying l'Hôpital's rule to the first of eqs. (20) we arrive at

$$\phi'(h) = \frac{1}{2g'(h)} \left[-U(\psi(h)) - Q^2 + Q^2 q^2 e^{4\phi(h)} \right] \quad (24)$$

which is negative as long as

$$U(\psi) \geq Q^2 (q^2 e^{4\phi} - 1) \quad (25)$$

at the horizon. For $U \geq 0$ we may read this as a condition on the electric charge,

$$|q| \leq e^{-2\phi(h)} \quad . \quad (26)$$

Looking back at eq. (19) we are able to complete the proof of the no-hair theorem in the traditional $\gamma=4$, $\epsilon=0$, $\frac{\partial V}{\partial \phi}=0$ case for arbitrarily small decay of $\psi \rightarrow \psi_0$, provided the conditions a)–c) are supplemented by e) an upper bound on the electric charge of the black hole.

In summary, we have written down all static classical solutions to a general class of two-dimensional dilaton-gravity-Maxwell theories and found them to generically exhibit multiple-horizon spacetimes. These solutions obey the classical no-hair theorem for scalar spectator fields. For the standard string effective spacetime action an improved argument does not rely on fall-off conditions for the spectator field but requires a bound on the electric charge of the black hole.

Acknowledgements: We acknowledge discussions with J. Schiff and E. Weinberg.

REFERENCES

1. R. Marnelius, *Nucl. Phys.* **B211** (1983), 14;
 C. Teitelboim, *Phys. Lett.* **B126** (1983), 41;
 R. Jackiw, in *Quantum Theory of Gravity*, S. Christensen, ed. (Hilger, Bristol U.K. 1984);
 A.H. Chamseddine, *Phys. Lett.* **B256** (1991), 379;
 T. Banks and M. O’Loughlin, *Nucl. Phys.* **B362** (1991), 649;
 R.B. Mann, “*Lower dimensional gravity*”, Waterloo preprint WATPHYS-TH-91-07 (Oct. 1991), to appear in Proc. of 4th Canadian Conf. on General Relativity and Relativistic Astrophysics, Winnipeg, Canada, May 1991.
2. E. Witten, *Phys. Rev.* **D44** (1991), 314;
 S. Elitzur, A. Forge and E. Rabinovici, *Nucl. Phys.* **B359** (1991), 581;
 G. Mandal, A. Sengupta and S. Wadia, *Mod. Phys. Lett.* **A6** (1991), 1685.
3. M.D. McGuigan, C.R. Nappi and S.A. Yost, “*Charged Black Holes in Two-Dimensional String Theory*,” IAS preprint IASSNS-HEP-91/57 (Oct. 91) hep-th/9111038, to appear in *Nucl. Phys.* **B**.
4. J.G. Russo and A.A. Tseytlin, “*Scalar-Tensor Quantum Gravity in Two Dimensions*,” Stanford and Cambridge preprint SU-ITP-92-2 DAMTP-1-1992 (Jan. 1992) hep-th/9201021;
 T.T. Burwick and A.H. Chamseddine, “*Classical and Quantum Considerations of Two-dimensional Gravity*,” Zürich preprint ZU-TH-4/92 (March 92) hep-th/9204002.
5. C.G. Callan, D. Friedan, E.J. Martinec and M.J. Perry, *Nucl. Phys.* **B262** (1985), 593;
 E.S. Fradkin and A.A. Tseytlin, *Nucl. Phys.* **B261** (1985), 1.
6. D. Garfinkle, G.T. Horowitz and A. Strominger, *Phys. Rev.* **D43** (1991), 3140.
7. G.W. Gibbons and S.W. Hawking, *Phys. Rev.* **D15** (1976), 2738.

8. C.G. Callan, S.B. Giddings, J.A. Harvey and A. Strominger, *Phys. Rev.* **D45** (1992), 1005.
9. S.P. de Alwis, “*Comments on No-Hair Theorems and Stability of Blackholes*”,
Boulder preprint COLO-HEP-267 (Jan. 1992) hep-th/9201053.
10. J.E. Chase, *Comm. Math. Phys.* **19** (1970), 276;
J.D. Bekenstein, *Phys. Rev.* **D5** (1972), 1239, *ibid.* **D5** (1972), 2403;
C. Teitelboim, *Phys. Rev.* **D5** (1972), 2941.
11. H.A. Buchdahl, *Phys. Rev.* **115** (1959), 1325;
M. Cadoni, *Phys. Rev.* **D44** (1991), 1115.